4764 14 P

# WORKSHOP ON EARLY MARS: HOW WARM AND HOW WET?

(NASA-CR-193503) WORKSHOP ON EARLY MARS: HOW WARM AND HOW WET, PART 2? (Lunar and Planetary Inst.) 14 p

N94-17637

**Unclas** 

G3/91 0194104





# LPI Technical Report Number 93-03, Part 2

Lunar and Planetary Institute 3600 Bay Area Boulevard Houston TX 77058-1113 LPI/TR--93-03, Part 2

.

# **WORKSHOP ON**

**EARLY MARS: HOW WARM AND HOW WET?** 

Edited by

S. Squyres and J. Kasting

Held in Breckenridge, Colorado

July 26-28, 1993

Sponsored by
MSATT Study Group
Lunar and Planetary Institute

Lunar and Planetary Institute 3600 Bay Area Boulevard Houston TX 77058-1113

LPI Technical Report Number 93-03, Part 2 LPI/TR--93-03, Part 2

# Compiled in 1993 by LUNAR AND PLANETARY INSTITUTE

The Institute is operated by the University Space Research Association under Contract No. NASW-4574 with the National Aeronautics and Space Administration.

Material in this volume may be copied without restraint for library, abstract service, education, or personal research purposes; however, republication of any paper or portion thereof requires the written permission of the authors as well as the appropriate acknowledgment of this publication.

This report may be cited as

Squyres S. and Kasting J., eds. (1993) Workshop on Early Mars: How Warm and How Wet? LPI Tech. Rpt. 93-03, Part 2, Lunar and Planetary Institute, Houston. 23 pp.

This report is distributed by

ORDER DEPARTMENT Lunar and Planetary Institute 3600 Bay Area Boulevard Houston TX 77058-1113

Mail order requestors will be invoiced for the cost of shipping and handling.

# **Preface**

In 1992 the MSATT program conducted a workshop on modeling of the martian climate. At that workshop it became clear that a serious problem had arisen concerning the early climate of Mars. Based on the evidence for smallscale fluvial activity, the view had been widely held that early in its history Mars had a climate that was much warmer and wetter than today's. However, most plausible recent climate models have fallen far short of the warm temperatures often inferred from the geologic evidence. Moreover, recent geophysical work has suggested that early geothermal warming may also have played a significant role in allowing fluvial activity. In order to address the issue of just how warm and how wet early Mars was, a workshop was convened in July of 1993, in Breckenridge, Colorado. The results of the workshop are reported here.

# PRECEDING PAGE BLANK NOT FILMED

Contents	_
Summary of Technical Sessions	1
List of Workshop Participants	3

. •	, or example of the second of	
<b>V</b>		

# PRECEDING PAGE BLANK NOT FILMED

# **Program**

### Monday, July 26, 1993

8:00 a.m.

Registration

8:45-9:45 a.m.

# **OVERVIEW OF MARS' EARLY EVOLUTION**

8:45 a.m.

Early Mars: Warmed from Without or Within

S. W. Squyres\*

9:15 a.m.

Possible Solutions to the Problem of Channel Formation on Early Mars

J. F. Kasting\*

9:45 a.m.

Break

### 10:00 a.m.-12:30 p.m.

# **VOLATILE INVENTORIES AND ATMOSPHERIC EVOLUTION**

10:00 a.m.

Wet Inside and Out?: Constraints on Water in the Martian Mantle and on

Outgassed Water, Based on Melt Inclusions in SNC Meteorites

H. Y. McSween Jr.\* and R. P. Harvey

10:30 a.m.

Mars Atmospheric Loss and Isotopic Fractionation by Solar-Wind-induced

Sputtering and Photochemical Escape

B. M. Jakosky\*, R. O. Pepin, R. E. Johnson, and J. L. Fox

11:00 a.m.

Evolution of the Martian Atmosphere

R. O. Pepin\*

11:30 a.m.

Requirements for the Early Atmosphere of Mars from the Nitrogen Isotope Ratios

J. L. Fox\*

12:00 p.m.

Adjourn

### 7:00-9:00 p.m.

# CLIMATE/GEOCHEMICAL/PHOTOCHEMICAL MODELING

7:00 p.m.

A Model for the Evolution of CO<sub>2</sub> on Mars

R. M. Haberle\*, D. Tyler, C. P. McKay, and W. L. Davis

<sup>\*</sup> Indicates speaker

Property and the second second

Early Mars: The Inextricable Link Between Internal and External Influences 7:30 p.m. on Valley Network Formation S. E. Postawko\* and F. P. Fanale The Young Sun and Photochemistry of the Primitive Martian Atmosphere 8:00 p.m. H. Nair\*, M. F. Gerstell, and Y. L. Yung A Carbon Dioxide/Methane Greenhouse Atmosphere on Early Mars 8:30 p.m. L. L. Brown\* and J. F. Kasting Mars and the Early Sun 9:00 p.m. D. Whitmire\*, L. R. Doyle, R. T. Reynolds, and P. G. Whitman Adjourn 9:30 p.m.

Tuesday, July 27, 1993 9:00 a.m.-12:30 p.m.

	MARTIAN HYDROLOGY AND VALLEY SYSTEM FORMATION
9:00 a.m.	The Changes on Mars at the End of Heavy Bombardment M. H. Carr*
9:30 a.m.	Evolution of the Global Water Cycle on Mars: The Geological Evidence V. R. Baker* and V. C. Gulick
10:00 a.m.	Fluvial Valleys in the Heavily Cratered Terrains of Mars: Evidence for Paleoclimatic Change? V. C. Gulick* and V. R. Baker
10:30 a.m.	Break
11:00 a.m.	The Hydrologic Response of Mars to the Onset of a Colder Climate and to the Thermal Evolution of Its Early Crust S. M. Clifford*
11:30 a.m.	Briny Lakes on Early Mars? Terrestrial Intracrater Playas and Martian Candidates P. Lee*
12:00 p.m.	Adjourn

# 7:00-9:30 p.m.

#### CLUES FROM MARTIAN GEOMORPHOLOGY

Early Mars: A Regional Assessment of Denudation Chronology 7:00 p.m.

T. A. Maxwell\* and R. A. Craddock

The Early Martian Environment: Clues from the Cratered Highlands and the 7:30 p.m.

Precambrian Earth

R. A. Craddock\* and T. A. Maxwell

Ancient Martian Valley Genesis and Paleoclimatic Inference: The Present as a 8:00 p.m.

Key to the Past

G. R. Brakenridge\*

Mars: Noachian Hydrology by Its Statistics and Topology 8:30 p.m.

N. A. Cabrol\* and E. A. Grin

The Martian Valley Networks: Origin By Niveo-Fluvial Processes 9:00 p.m.

J. Rice\*

Adjourn 9:30 p.m.

Wednesday, July 28, 1993 9:00-10:00 a.m.

#### CHEMICAL WEATHERING AND CLIMATE

An Attempt to Comprehend Martian Weathering Conditions Through the Analysis 9:00 a.m.

of Terrestrial Palagonite Samples

C. Douglas\*, I. P. Wright, J. B. Bell, R. V. Morris, D. C. Golden,

and C. T. Pillinger

Oxidation of Dissolved Iron Under Warmer, Wetter Conditions on 9:30 a.m.

Mars: Transitions to Present-Day Arid Environments

R. G. Burns\*

10:00 a.m. Break

10:30 a.m. Discussion

Adjourn 12:30 p.m.

, , , , , , , , , , , , , , , , , , ,			
			·
			·
			·

# **Summary of Technical Sessions**

It is clear that conditions on Mars during the Noachian Period (which extended from the beginning of Mars' geologic record to something like 3.9 b.y. ago) were significantly different than they are today. As on the other planets, the rate of meteoritic bombardment was far higher than it is at present; this is shown clearly by the densely cratered nature of Noachian terrains. But there were other differences as well. Some are only inferred, while for others we have clear evidence. The nature of these differences, and what they imply about broader topics of planetary evolution, were the primary subjects of the workshop.

The atmosphere of Mars during the Noachian was probably substantially more dense than the present atmosphere of 7 mbar of CO<sub>2</sub>. Just how dense it was is a subject of some debate. Processes that fed it during this period include volcanic outgassing and infall of volatile-rich meteorites. Processes that destroyed it may have included hydrodynamic and thermal escape, solar wind erosion, impact erosion, and chemical processes at the surface such as carbonate formation. The balance among these various processes has been a topic of much recent work and was vigorously discussed at the workshop. While the atmospheric pressure during the Noachian is unknown, there is a consensus that pressures of up to a few bars are plausible. If this was the case, then carbonate formation during or after the Noachian must have been important. Other sinks could remove hundreds of millibars of CO<sub>2</sub> from the atmosphere, but removal of a few bars would require that a substantial carbonate reservoir currently be present on Mars.

Probably another important characteristic of the Noachian was a much higher geothermal heat flow. Detailed models of the accretion and early thermal evolution of Mars, as well as isotopic evidence from SNC meteorites believed to come from Mars, provide persuasive evidence for very early core formation, a hot early mantle, and a high early heat flow. This recognition that Mars probably began its history hot has been one of the major developments in martian geophysics over the last several years. The mean global heat flow during the Noachian is also very uncertain, but a value 3.9 b.y. ago of about 5 × the present one (which is typically estimated at about 45-50 mW/m<sup>2</sup>) is plausible. Whatever the global mean heat flow was at that time, it is certain that local values in areas of enhanced geothermal activity were significantly higher still.

Given these very different environmental conditions, it is natural to expect that geomorphic processes during the Noachian were different as well, and it is clear that they were. The most obvious difference is that the rate of formation of valley systems was far higher during the Noachian than it has been during any other period of martian history. Valley systems are small, quasidendritic drainage networks found primarily in the ancient cratered highlands of Mars. Typical valley widths are 1-2 km, and lengths extend up to several hundred kilometers. The valleys were almost certainly carved by liquid water flowing at low discharges in small channels. These valley systems have long provided the primary physical evidence suggesting that conditions on Mars were different early in the planet's history.

There is other morphologic evidence as well. Careful comparison of the depths and rim heights of Noachian craters with post-Noachian craters shows clearly that the rate of crater degradation (lowering of crater rims and raising of crater floors) was far higher in the Noachian than it has been since then. Such degradation is certainly attributable in part to erosion and deposition, though other processes (such as volcanic infilling) could also have played a role.

There is general agreement that there has been enough H<sub>2</sub>O on Mars for us to accept that water was the fluid that carved the valley systems. However, the details of their formation are unclear. In particular, there are two key obviously linked questions that remain unresolved: Did conditions during the Noachian allow precipitation? And does the geomorphic evidence from the Noachian require precipitation?

If the climate of Mars was warm enough during the Noachian to allow precipitation, then an enhanced greenhouse effect enabled by a denser atmosphere is required. However, temperatures anywhere close to the melting point of water may be difficult to achieve with a dense early CO<sub>2</sub> atmosphere on Mars. Earlier models predicted substantial greenhouse warming for an atmosphere of a few bars of CO<sub>2</sub>. However, more recent models that include the effects of CO<sub>2</sub> condensation and cloud formation suggest that such an atmosphere is capable of raising the martian surface temperature by at most several tens of degrees. For the conventional view of solar evolution (which suggests a luminosity about 75% the present value during the Noachian), such a greenhouse does not appear capable of producing mean surface temperatures in excess of ~230 K.

There are two other factors that could conceivably contribute further climatic warming, but both are highly uncertain. One is the possibility that the early Sun was substantially more luminous than predicted by conventional models; this would require a much higher rate of solar mass loss during the Sun's early history than is usually accepted. The other is the possibility that substantial quantities of methane, another very effective greenhouse gas, were present in the martian atmosphere. Without either of these effects, however, the best recent work suggests that climatic warming to temperatures above 273 K during the Noachian was not possible.

There also is no general agreement as to whether the geomorphic evidence actually requires precipitation. Some researchers point to the ubiquitous nature of the valley systems in old heavily cratered terrain on Mars, and to the fact that some valley heads extend right to the crests of drainage divides. From these characteristics they infer that precipitation and runoff must have taken place. Others note that the morphology of many valley systems is more consistent with sapping than runoff, and that the highest drainage densities on Mars are substantially lower than those in even arid regions on Earth. From these characteristics, they infer that the valleys formed by sapping of subsurface liquid, and that precipitation was not required. There is also disagreement as to whether the early high rates of crater degradation require precipitation, or whether they are just a consequence of higher early rates of sapping, eolian transport, volcanism, and so forth.

There are geomorphic points on which there is agreement. There are some valley systems for which all workers agree that sapping is a satisfactory explanation, without the need to invoke precipitation and runoff. It is also clear that many valley systems are too extensive to have formed by the single discharge of an aquifer, i.e., that some type of aquifer recharge is required. Several different processes, including precipitation and hydrothermal convection (enabled by the early high heat flow), have been examined closely as recharge mechanisms. Both of these appear to be more than adequate to do the job. The basic difficulties are that dissimilar processes can produce similar final landforms, and that many

processes may have operated on these valleys, including ones that have substantially obscured the evidence for the primary genetic process.

A final issue that was considered at the workshop was identification of directions for future work. Important areas where further progress should be possible include these:

- Use geochemical data from Mars Observer or a follow-on mission to search for martian carbonates.
- Use imaging data from Mars Observer or a followon mission to search for further morphologic evidence requiring Noachian precipitation.
- Improve models of the contribution of methane to the martian greenhouse.
- Further investigate the prospects for a more luminous early Sun.
- Model the climatic effects of the very high and rapidly varying obliquities that are suggested by recent dynamical modeling.
- Improve climate models to include the optical properties of CO<sub>2</sub> clouds.
- Continue investigation of terrestrial analog terrains and processes.

Note added in proof: In light of the failure of Mars Observer, a top priority should be a reflight of the instruments to obtain the data that otherwise will not be acquired.

# **List of Workshop Participants**

#### J. R. Baker

Department of Geosciences University of Arizona Tucson AZ 85721 Phone: 602-621-6003 Fax: 602-621-2672

#### J. Bergstralh

Code SLC

NASA Headquarters Washington DC 20546 Phone: 202-453-1580 Fax: 202-358-3097

E-mail: NHQVAX::JBERGSTR

#### Robert Brakenridge

Surficial Processes Laboratory Department of Geography Dartmouth College Hanover NH 03755 Phone: 603-646-2870 Fax: 603-646-2810

E-mail: brakenridge@dartmouth.edu

#### Lisa L. Brown

Department of Geosciences 332 Deike Building Pennsylvania State University University Park PA 16802 Phone: 814-863-3965 Fax: 814-865-3191

E-mail: brown@essc.psu.edu

#### Mark Bullock

Laboratory for Atmospheric and Space Physics

Campus Box 392 University of Colorado Boulder CO 80309-0392 Phone: 303-492-1628 Fax: 303-492-6946

E-mail: ZODIAC::BULLOCK

#### Roger G. Burns

Mail Code 54-816

Massachusetts Institute of Technology

Cambridge MA 02139 Phone: 617-253-1906 Fax: 617-253-6208

E-mail: bcabral@eagle.mit.edu

#### Bryan Butler

Division of Geological and Planetary Science Mail Stop 170-25

California Institute of Technology

Pasadena CA 91125 Phone: 818-356-6477 Fax: 818-585-1917

E-mail: butler@scat.gps.caltech.edu

#### Nathalie Cabrol

Telescope de 1 Metre Observatoire de Paris 92195 Meudon Cedex Principal **FRANCE** 

Phone: 45-07-75-30 Fax: 45-07-79-59

E-mail: MESIOA::DOLLFUS

#### Michael H. Carr

Mail Stop 946 U.S. Geological Survey 345 Middlefield Road Menlo Park CA 94025 Phone: 415-329-5174 Fax: 415-329-4936

E-mail: mcarr@isdmnl.wr.usgs.gov

### Michael Carroll

Sky and Telescope Magazine 6820 W. Chestnut Avenue Littleton CO 80123 Phone: 303-933-1645

#### Benton Clark

Mail Stop B0560 Martin Marietta P.O. Box 179 Denver CO 80201 Phone: 303-971-9007 Fax: 303-971-9141

E-mail: bclark@den.mmc.com

#### Stephen Clifford

Lunar and Planetary Institute 3600 Bay Area Boulevard Houston TX 77058 Phone: 713-486-2146 Fax: 713-486-2162

E-mail: clifford@lpi.jsc.nasa.gov

#### Robert A. Craddock

Center for Earth and Planetary Studies National Air and Space Museum, Room 3775

MRC 315

Smithsonian Institution Washington DC 20560 Phone: 202-357-1457 Fax: 202-786-2566

E-mail: craddock@ceps.edu

#### David DesMarais

Mail Stop 239-4

NASA Ames Research Center Moffett Field CA 94035-1000 Phone: 415-604-3220

Fax: 415-604-1088

E-mail: david\_desmarais@qmgate.arc.nasa.gov

#### Tammy Dickinson

Code SLC

NASA Headquarters

Washington DC 20546-0001 Phone: 202-358-0292 Fax: 202-358-3097

E-mail: tdickinson@nasamail.nasa.gov

#### Caroline Douglas

Planetary Science Unit
Department of Earth Sciences

Walton Hall Open University

Milton Keynes MK7 6AA UNITED KINGDOM Phone: 0908-655-988

Fax: 0908-655-151

#### Gerlind Dreibus

Max-Planck-Institut für Chemie

Saarstrasse 23 D-6500 Mainz GERMANY

Phone: 49-6131-305-395 Fax: 49-6131-371-290

#### Jane L. Fox

**MSRC** 

Institute for Terrestrial and Planetary Atmospheres

State University of New York Stony Brook NY 11794 Phone: 516-632-8317 Fax: 516-632-8820

E-mail: fox@atmsci.sunysb.edu

#### Virginia Gulick

Mail Stop 245-3

Space Science Division NASA Ames Research Center Moffett Field CA 94035

Phone: 415-604-0781 Fax: 415-604-6779

E-mail: gulick@convx1.ccit.arizona.edu

#### R. M. Haberle

Mail Stop 245-3

Space Science Division NASA Ames Research Center Moffett Field CA 94035 Phone: 415-604-5491

Fax: 415-604-6779

E-mail: haberle@humbabe.arc.nasa.gov

#### Joan Hayashi

Laboratory for Atmospheric and Space Physics

Campus Box 392 University of Colorado Boulder CO 80309 Phone: 303-492-8918 Fax: 303-492-6946

E-mail: hayashi@argyre.colorado.edu

#### Bruce Jakosky

Laboratory for Atmospheric and Space Physics

Campus Box 392 University of Colorado Boulder CO 80309-0392 Phone: 303-492-8004 Fax: 303-492-6946

E-mail: ZODIAC::JAKOSKY

#### John Jones

Mail Code SN2

NASA Johnson Space Center

Houston TX 77058 Phone: 713-483-5319 Fax: 713-483-5347 E-mail: SN::JJONES

#### James Kasting

211 Deike

Pennsylvania State University University Park PA 16802 Phone: 814-865-3207 Fax: 814-865-3191

E-mail: kasting@essc.psu.edu

#### Pascal Lee

404 Space Sciences Building Cornell University Ithaca NY 14853-6801

Phone: 607-255-6237 Fax: 607-255-9002

E-mail: lee@astronsun.tn.cornell.edu

#### Ted A. Maxwell

National Air and Space Museum

MRC 315

Smithsonian Institution Washington DC 20560 Phone: 202-357-1424 Fax: 202-786-2566

E-mail: tmaxwell@nasamail.nasa.gov

#### Daniel J. McCleese

Mail Stop 183-335 Jet Propulsion Laboratory 4800 Oak Grove Drive Pasadena CA 91109 Phone: 818-354-2317

Fax: 818-393-6546

E-mail: djmcc@scn1.jpl.nasa.gov

#### Harry McSween

Department of Geological Sciences

University of Tennessee Knoxville TN 37996-1410 Phone: 615-974-5498 Fax: 615-974-2368

E-mail: mcsween@utkvx.utk.edu

#### Mike Mellon

Laboratory for Atmospheric and Space Physics

Campus Box 392 University of Colorado Boulder CO 80302 Phone: 303-492-7902 Fax: 303-492-6946

E-mail: mellon@argyre.colorado.edu

#### Michael A. Meyer

Lockheed Engineering and Sciences Co.

500 E Street, SW Suite 800

Washington DC 20024 Phone: 202-863-5257 Fax: 202-863-8217

E-mail: mmyer@nasamail.nasa.gov

#### Hari A. Nair

Mail Stop 170-25

California Institute of Technology

Pasadena CA 91125 Phone: 818-356-6960 Fax: 818-585-1917

E-mail: hari@mercu1.gps.caltech.edu

#### Robert O. Pepin

School of Physics and Astronomy University of Minnesota 116 Church Street, SE Minneapolis MN 55455

Phone: 612-624-0819 Fax: 612-624-4578

#### Jeff Plescia

Mail Stop 183-501 Jet Propulsion Laboratory 4800 Oak Grove Drive

Pasadena CA 91109 Phone: 818-354-6936 Fax: 818-354-0966

E-mail: iplescia@nasamail.nasa.gov

#### Susan Postawko

School of Meteorology University of Oklahoma 100 East Boyd Norman OK 73019 Phone: 405-325-6561

Fax: 405-325-7689

E-mail: spostawko@geohub.gen.uoknor.edu

Department of Geography Arizona State University Tempe AZ 85287 Phone: 602-496-6512 Fax: 602-965-8313

E-mail: asjwr@asuvm.inre.asu.edu

#### Ben Schuraytz

Lunar and Planetary Institute 3600 Bay Area Boulevard Houston TX 77058 Phone: 713-486-2187

Fax: 713-486-2162

E-mail: schuraytz@lpi.jsc.nasa.gov

#### Bernhard Spettel

Abteiling Kosmochemie Max-Planck-Institut für Chemie Saarstrasse 23 D-6500 Mainz **GERMANY** 

Phone: 49-6131-305-233

#### Steve Squyres

Space Sciences Building Cornell University Ithaca NY 14853 Phone: 607-255-3508

Fax: 607-255-5907

E-mail: squyres@astrosun.tn.cornell.edu

#### John Stansberry

Department of Planetary Sciences

University of Arizona Tucson AZ 85721 Phone: 602-621-2276

E-mail: stansber@lpl.arizona.edu

#### Dan Tyler Jr.

2561 Mardell Way Mountain View CA 94043 Phone: 415-604-0322

E-mail: tyler@oskar.ames.nasa.gov

#### Daniel P. Whitmire

Department of Physics 44210 University of Southwestern Louisiana

Phone: 318-231-6185 Fax: 318-321-6190 E-mail: dpw9254@usl.edu

Lafayette LA 70504

#### Frank Wlotzka

Abteilung Kosmochemie Max-Planck-Institut für Chemie Saarstrasse 23 D-6500 Mainz **GERMANY** 

Phone: 49-6131-305-343 Fax: 49-6131-371-290

E-mail: włotzka@mpch-mainz.mpg.dbp.de

#### Aaron Zent

Mail Stop 245-3 NASA Ames Research Center Moffett Field CA 94035

Phone: 415-604-5517 Fax: 415-604-6779

E-mail: zent@pan.arc.nasa.gov

#### Robert Zubrin

Martin Marietta P.O. Box 179 Denver CO 80201 Phone: 303-971-9299

Fax: 303-971-3600